



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

by a modicum of the new. In fact, the author seems to regard modern morphologists as perhaps too narrow in their vision to possess the proper perspective for a general text. As a result, the two volumes now published are distinctly not up to date from the standpoint of modern morphology.

The first volume (pp. 277), issued in 1905, is introduced by general discussions of questions that are not current in modern laboratories, however interesting they may have been during the domination of the Goethean morphology. The main part of the volume consists of "Die Morphologie der Kryptogamen." The illustrations are well selected and many of them are original. The ground has evidently been well traversed in the laboratory but not in recent literature, for the text gives a distinct impression of long isolation. Why do so many botanists fail to see the incongruity of such a series of names for coordinate groups as "A. Thallophyten," "B. Charophyta," "C. Moose," "D. Gefässkryptogamen"?

The second volume (pp. 454), issued in 1907, presents "Die Morphologie der Phanerogamen," in so far as the vegetative organs are concerned. The main divisions are "Die Keimpflanze," "Die Wurzel," "Das Blatt," "Die Achse," and "Die Trichome." The general attitude is indicated by the subdivisions under "Seedlings," which are as follows: seedlings of polycotyls, of monocotyls, of stemless plants, of acotylous plants.

The third volume is to contain the morphology of the "phanerogam flower" and a system of classification, besides a general index.

The work will undoubtedly prove a most interesting one for reference, especially for the younger generation of morphologists.—J. M. C.

MINOR NOTICES

North American Flora.—The first part of volume xxv has just appeared, being the fifth part published. It contains four of the sixteen families of Geraniales, the order with a synopsis of its families being presented by J. K. SMALL. The Geraniaceae, by L. T. HANKS and J. K. SMALL, contain four genera, a new genus *Robertiella* being founded on *Geranium Robertianum*. The species of *Geranium* number sixty-four, of which twenty-three are new, and *Erodium* and *Pelargonium* each contain six species. The Oxalidaceae, by J. K. SMALL, contain ten genera, *Hesperoxalis*, *Otoxalis*, and *Bolboxalis* being new and monotypic. The other genera are *Oxalis* (3 spp., 1 new), *Ionoxalis* (65 spp., 28 new), *Monoxalis* (2 spp., 1 new), *Lotoxalis* (11 spp., 1 new), *Xanthoxalis* (26 spp., 1 new), *Biophytum* (1 sp.), and *Averrhoa* (2 spp.). The Erythroxylaceae, by N. L. BRITTON, represented by the single genus *Erythroxylon*, contains twenty-eight species, one of which is new. The Linaceae, by J. K. SMALL, contain four genera, *Hesperolinon* being a new genus established on *Linum* § *Hesperolinon* Gray. The genera are *Linum* (5 spp.), *Cathartolinum* (48 spp., 10 new and all but one of the remainder transferred from *Linum* by SMALL), *Hesperolinon* (9 spp.), and *Reinwardtia* (1 sp.).—J. M. C.